

## INTRODUCTION

5

10

15

## 25

30

09811-0810

09811-0810

09811-0810

09811-0810

improved by detecting new markers in cells, tissues, or bodily fluids which could differentiate between different stages of invasion.

Accordingly, there is a great need for more sensitive  
5 and accurate methods for the staging of breast cancer in a human to determine whether or not such cancer has metastasized and for monitoring the progress of breast cancer in a human which has not metastasized for the onset of metastasis.

In the present invention, methods are provided for  
10 detecting, diagnosing, monitoring, staging, prognosticating, imaging and treating breast cancer via breast specific genes referred to herein as BSGs. For purposes of the present invention, BSG refers, among other things, to native protein expressed by the gene comprising a polynucleotide sequence of  
15 SEQ ID NO:1, 2, 3, 4 or 5. By "BSG" it is also meant herein polynucleotides which, due to degeneracy in genetic coding, comprise variations in nucleotide sequence as compared to SEQ ID NO: 1, 2, 3, 4 or 5 but which still encode the same protein. In the alternative, what is meant by BSG as used  
20 herein, means the native mRNA encoded by the gene comprising the polynucleotide sequence of SEQ ID NO: 1, 2, 3, 4 or 5, levels of the gene comprising the polynucleotide sequence of SEQ ID NO: 1, 2, 3, 4 or 5, or levels of a polynucleotide which is capable of hybridizing under stringent conditions to  
25 the antisense sequence of SEQ ID NO: 1, 2, 3, 4 or 5. An exemplary BSG polypeptide encoded by the BSG polynucleotide of SEQ ID NO: 4 is depicted in SEQ ID NO:6.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in  
30 the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention are given by way of illustration only. Various changes and modifications within the spirit and scope of the  
35 disclosed invention will become readily apparent to those

09886241.062101  
FOI290729866

skilled in the art from reading the following description and from reading the other parts of the present disclosure.

#### SUMMARY OF THE INVENTION

5           Toward these ends, and others, it is an object of the present invention to provide BSGs comprising a polynucleotide of SEQ ID NO: 1, 2, 3, 4, or 5, a protein expressed by a polynucleotide of SEQ ID NO: 1, 2, 3, 4, or 5, such as that of SEQ ID NO:6, or a variant thereof which expresses the  
10 protein; or a polynucleotide which is capable of hybridizing under stringent conditions to the antisense sequence of SEQ ID NO:1, 2, 3, 4, or 5.

It is another object of the present invention to provide a method for diagnosing the presence of breast cancer by  
15 analyzing for changes in levels of BSG in cells, tissues or bodily fluids compared with levels of BSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein a change in levels of BSG in the patient versus the normal human control is associated with breast  
20 cancer.

Further provided is a method of diagnosing metastatic breast cancer in a patient having breast cancer which is not known to have metastasized by identifying a human patient suspected of having breast cancer that has metastasized;  
25 analyzing a sample of cells, tissues, or bodily fluid from such patient for BSG; comparing the BSG levels in such cells, tissues, or bodily fluid with levels of BSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in BSG levels in the patient  
30 versus the normal human control is associated with breast cancer which has metastasized.

Also provided by the invention is a method of staging breast cancer in a human which has such cancer by identifying a human patient having such cancer; analyzing a sample of  
35 cells, tissues, or bodily fluid from such patient for BSG;

0936241 03101  
FOR 290 " 429350

comparing BSG levels in such cells, tissues, or bodily fluid with levels of BSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in BSG levels in the patient versus the normal  
5 human control is associated with a cancer which is progressing and a decrease in the levels of BSG is associated with a cancer which is regressing or in remission.

Further provided is a method of monitoring breast cancer in a human having such cancer for the onset of metastasis.  
10 The method comprises identifying a human patient having such cancer that is not known to have metastasized; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for BSG; comparing the BSG levels in such cells, tissue, or bodily fluid with levels of BSG in preferably the  
15 same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in BSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.

Further provided is a method of monitoring the change  
20 in stage of breast cancer in a human having such cancer by looking at levels of BSG in a human having such cancer. The method comprises identifying a human patient having such cancer; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for BSG; comparing the BSG  
25 levels in such cells, tissue, or bodily fluid with levels of BSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in BSG levels in the patient versus the normal human control is associated with a cancer which is progressing and a decrease  
30 in the levels of BSG is associated with a cancer which is regressing or in remission.

Further provided are methods of designing new therapeutic agents targeted to a BSG for use in imaging and treating breast cancer. For example, in one embodiment,  
35 therapeutic agents such as antibodies targeted against BSG or

09886241-062101

fragments of such antibodies can be used to treat, detect or image localization of BSG in a patient for the purpose of detecting or diagnosing a disease or condition. In this embodiment, an increase in the amount of labeled antibody  
5 detected as compared to normal tissue would be indicative of tumor metastases or growth. Such antibodies can be polyclonal, monoclonal, or omniclonal or prepared by molecular biology techniques. The term "antibody", as used herein and throughout the instant specification is also meant to include  
10 aptamers and single-stranded oligonucleotides such as those derived from an *in vitro* evolution protocol referred to as SELEX and well known to those skilled in the art. Antibodies can be labeled with a variety of detectable and therapeutic labels including, but not limited to, radioisotopes and  
15 paramagnetic metals. Therapeutic agents such as small molecules and antibodies which decrease the concentration and/or activity of BSG can also be used in the treatment of diseases characterized by overexpression of BSG. Such agents can be readily identified in accordance with teachings herein.  
20 Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of  
25 the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

30

#### GLOSSARY

The following illustrative explanations are provided to facilitate understanding of certain terms used frequently herein, particularly in the examples. The explanations are

09836241 "062101"

provided as a convenience and are not limitative of the invention.

ISOLATED means altered "by the hand of man" from its natural state; i.e., that, if it occurs in nature, it has been  
5 changed or removed from its original environment, or both.

For example, a naturally occurring polynucleotide or a polypeptide naturally present in a living animal in its natural state is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its  
10 natural state is "isolated", as the term is employed herein. For example, with respect to polynucleotides, the term isolated means that it is separated from the chromosome and cell in which it naturally occurs.

As part of or following isolation, such polynucleotides  
15 can be joined to other polynucleotides, such as DNAs, for mutagenesis, to form fusion proteins, and for propagation or expression in a host, for instance. The isolated polynucleotides, alone or joined to other polynucleotides such as vectors, can be introduced into host cells, in culture or  
20 in whole organisms. When introduced into host cells in culture or in whole organisms, such DNAs still would be isolated, as the term is used herein, because they would not be in their naturally occurring form or environment. Similarly, the polynucleotides and polypeptides may occur in a composition,  
25 such as media formulations, solutions for introduction of polynucleotides or polypeptides, for example, into cells, compositions or solutions for chemical or enzymatic reactions, for instance, which are not naturally occurring compositions, and, therein remain isolated polynucleotides or polypeptides  
30 within the meaning of that term as it is employed herein.

OLIGONUCLEOTIDE(S) refers to relatively short polynucleotides. Often the term refers to single-stranded deoxyribonucleotides, but it can refer as well to single-or double-stranded ribonucleotides, RNA:DNA hybrids and double-  
35 stranded DNAs, among others.

05386241-062101

Oligonucleotides, such as single-stranded DNA probe oligonucleotides, often are synthesized by chemical methods, such as those implemented on automated oligonucleotide synthesizers. However, oligonucleotides can be made by a  
5 variety of other methods, including *in vitro* recombinant DNA-mediated techniques and by expression of DNAs in cells and organisms.

Initially, chemically synthesized DNAs typically are obtained without a 5' phosphate. The 5' ends of such  
10 oligonucleotides are not substrates for phosphodiester bond formation by ligation reactions that employ DNA ligases typically used to form recombinant DNA molecules. Where ligation of such oligonucleotides is desired, a phosphate can be added by standard techniques, such as those that employ a  
15 kinase and ATP.

The 3' end of a chemically synthesized oligonucleotide generally has a free hydroxyl group and, in the presence of a ligase such as T4 DNA ligase, readily will form a phosphodiester bond with a 5' phosphate of another  
20 polynucleotide, such as another oligonucleotide. As is well known, this reaction can be prevented selectively, where desired, by removing the 5' phosphates of the other polynucleotide(s) prior to ligation.

POLYNUCLEOTIDE(S) generally refers to any  
25 polyribonucleotide or polydeoxribonucleotide and is inclusive of unmodified RNA or DNA as well as modified RNA or DNA. Thus, for instance, polynucleotides as used herein refers to, among other things, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single-  
30 and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, polynucleotide, as used herein, refers  
35 to triple-stranded regions comprising RNA or DNA or both RNA

09886241.062101  
1012290.14298860



and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the 5 molecules of a triple-helical region often is an oligonucleotide.

As used herein, the term polynucleotide is also inclusive of DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones 10 modified for stability or for other reasons are "polynucleotides" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein.

15 It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term polynucleotide as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of 20 polynucleotides, as well as chemical forms of DNA and RNA characteristic of viruses and cells, including simple and complex cells, inter alia.

POLYPEPTIDES, as used herein, includes all polypeptides as described below. The basic structure of polypeptides is 25 well known and has been described in innumerable textbooks and other publications in the art. In this context, the term is used herein to refer to any peptide or protein comprising two or more amino acids joined to each other in a linear chain by peptide bonds. As used herein, the term refers to both short 30 chains, which also commonly are referred to in the art as peptides, oligopeptides and oligomers, for example, and to longer chains, which generally are referred to in the art as proteins, of which there are many types. It will be appreciated that polypeptides often contain amino acids other 35 than the 20 amino acids commonly referred to as the 20

09885241.062101

naturally occurring amino acids, and that many amino acids, including the terminal amino acids, may be modified in a given polypeptide, either by natural processes such as processing and other post-translational modifications, or by chemical  
5 modification techniques which are well known to the art. Even the common modifications that occur naturally in polypeptides are too numerous to list exhaustively here, but they are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well  
10 known to those of skill in the art.

Modifications which may be present in polypeptides of the present invention include, to name an illustrative few, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety,  
15 covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine,  
20 formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated  
25 addition of amino acids to proteins such as arginylation, and ubiquitination.

Such modifications are well known to those of skill and have been described in great detail in the scientific literature. Several particularly common modifications  
30 including, but not limited to, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation are described in most basic texts, such as, for instance PROTEINS STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman  
35 and Company, New York (1993). Many detailed reviews are

09886241-062101



polypeptide amino acid sequence. For instance, as is well known, glycosylation often does not occur in bacterial hosts such as *E. coli*. Accordingly, when glycosylation is desired, a polypeptide can be expressed in a glycosylating host, generally a eukaryotic cell. Insect cells often carry out the same posttranslational glycosylations as mammalian cells. Thus, insect cell expression systems have been developed to express efficiently mammalian proteins having native patterns of glycosylation, *inter alia*. Similar considerations apply to other modifications.

It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications.

In general, as used herein, the term polypeptide encompasses all such modifications, particularly those that are present in polypeptides synthesized by expressing a polynucleotide in a host cell.

VARIANT(S) of polynucleotides or polypeptides, as the term is used herein, are polynucleotides or polypeptides that differ from a reference polynucleotide or polypeptide, respectively.

With respect to variant polynucleotides, differences are generally limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many regions, identical. Thus, changes in the nucleotide sequence of the variant may be silent. That is, they may not alter the amino acids encoded by the polynucleotide. Where alterations are limited to silent changes of this type a variant will encode a polypeptide with the same amino acid sequence as the reference. Alternatively, changes in the nucleotide sequence of the variant may alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Such nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and

09386241-062101

truncations in the polypeptide encoded by the reference sequence.

With respect to variant polypeptides, differences are generally limited so that the sequences of the reference and the variant are closely similar overall and, in many region, identical. For example, a variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions, fusions and truncations, which may be present in any combination.

10 RECEPTOR MOLECULE, as used herein, refers to molecules which bind or interact specifically with BSG polypeptides of the present invention and is inclusive not only of classic receptors, which are preferred, but also other molecules that specifically bind to or interact with polypeptides of the  
15 invention (which also may be referred to as "binding molecules" and "interaction molecules," respectively and as "BSG binding or interaction molecules". Binding between polypeptides of the invention and such molecules, including receptor or binding or interaction molecules may be exclusive  
20 to polypeptides of the invention, which is very highly preferred, or it may be highly specific for polypeptides of the invention, which is highly preferred, or it may be highly specific to a group of proteins that includes polypeptides of the invention, which is preferred, or it may be specific to  
25 several groups of proteins at least one of which includes polypeptides of the invention.

Receptors also may be non-naturally occurring, such as antibodies and antibody-derived reagents that bind to polypeptides of the invention.

30

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to novel breast specific polypeptides and polynucleotides, referred to herein as BSGs, among other things, as described in greater detail below.

35 ***Polynucleotides***

09836244-062101

In accordance with one aspect of the present invention, there are provided isolated BSG polynucleotides which encode BSG polypeptides.

Using the information provided herein, such as the  
5 polynucleotide sequences set out in SEQ ID NO:1, 2, 3, 4, or 5, a polynucleotide of the present invention encoding a BSG may be obtained using standard cloning and screening procedures, such as those for cloning cDNAs using mRNA from cells of a human tumor as starting material.

10 Polynucleotides of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The DNA may be double-stranded or single-stranded.  
15 Single-stranded DNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

The coding sequence which encodes the polypeptides may be identical to the coding sequence of the polynucleotides of  
20 SEQ ID NO:1, 2, 3, 4, or 5. It also may be a polynucleotide with a different sequence, which, as a result of the redundancy (degeneracy) of the genetic code, encodes the same polypeptides as encoded by SEQ ID NO:1, 2, 3, 4, or 5.

Polynucleotides of the present invention, such as SEQ  
25 ID NO: 1, 2, 3, 4, or 5, which encode these polypeptides may comprise the coding sequence for the mature polypeptide by itself. Polynucleotides of the present invention may also comprise the coding sequence for the mature polypeptide and additional coding sequences such as those encoding a leader  
30 or secretory sequence such as a pre-, or pro- or prepro-protein sequence. Polynucleotides of the present invention may also comprise the coding sequence of the mature polypeptide, with or without the aforementioned additional coding sequences, together with additional, non-coding  
35 sequences. Examples of additional non-coding sequences which

09886241-062101

may be incorporated into the polynucleotide of the present invention include, but are not limited to, introns and non-coding 5' and 3' sequences such as transcribed, non-translated sequences that play a role in transcription, mRNA processing including, for example, splicing and polyadenylation signals, ribosome binding and stability of mRNA, and additional coding sequence which codes for amino acids such as those which provide additional functionalities. Thus, for instance, the polypeptide may be fused to a marker sequence such as a peptide which facilitates purification of the fused polypeptide. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, such as the tag provided in the pQE vector (Qiagen, Inc.), among others, many of which are commercially available. As described in Gentz et al. (Proc. Natl. Acad. Sci., USA 86: 821-824 (1989)), for instance, hexa-histidine provides for convenient purification of the fusion protein. The HA tag corresponds to an epitope derived of influenza hemagglutinin protein (Wilson et al., Cell 37: 767 (1984)).

In accordance with the foregoing, the term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides which include a sequence encoding a polypeptide of the present invention, particularly SEQ ID NO:1, 2, 3, 4, or 5. The term encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, interrupted by introns) together with additional regions, that also may contain coding and/or non-coding sequences.

The present invention further relates to variants of the herein above described polynucleotides which encode for fragments, analogs and derivatives of the BSG polypeptides. A variant of the polynucleotide may be a naturally occurring variant such as a naturally occurring allelic variant, or it may be a variant that is not known to occur naturally. Such non-naturally occurring variants of the polynucleotide may be

made by mutagenesis techniques, including those applied to polynucleotides, cells or organisms.

Among variants in this regard are variants that differ from the aforementioned polynucleotides by nucleotide  
5 substitutions, deletions or additions. The substitutions, deletions or additions may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions,  
10 deletions or additions.

Among the particularly preferred embodiments of the invention in this regard are polynucleotides encoding polypeptides having the same amino acid sequence encoded by a BSG polynucleotide comprising SEQ ID NO: 1, 2, 3, 4, or 5; 15 variants, analogs, derivatives and fragments thereof, and fragments of the variants, analogs and derivatives. Further particularly preferred in this regard are BSG polynucleotides encoding polypeptide variants, analogs, derivatives and fragments, and variants, analogs and derivatives of the 20 fragments, in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of the BSG. Also 25 especially preferred in this regard are conservative substitutions. Most highly preferred are polynucleotides encoding polypeptides having the amino acid sequences as polypeptides encoded by SEQ ID NO: 1, 2, 3, 4, or 5, without substitutions.

Further preferred embodiments of the invention are BSG polynucleotides that are at least 70% identical to a polynucleotide of SEQ ID NO: 1, 2, 3, 4, or 5, and polynucleotides which are complementary to such polynucleotides. More preferred are BSG polynucleotides that comprise a region that is at least 80% identical to a



polynucleotide of SEQ ID NO: 1, 2, 3, 4, or 5. In this regard, BSG polynucleotides at least 90% identical to the same are particularly preferred, and among these particularly preferred BSG polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the most preferred.

10 Particularly preferred embodiments in this respect, moreover, are polynucleotides which encode polypeptides which retain substantially the same biological function or activity as the mature polypeptides encoded by a polynucleotide of SEQ ID NO: 1, 2, 3, 4, or 5.

15 The present invention further relates to polynucleotides that hybridize to the herein above-described BSG sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, 20 the term "stringent conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences.

As discussed additionally herein regarding polynucleotide assays of the invention, for instance, 25 polynucleotides of the invention as described herein, may be used as a hybridization probe for cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding BSGs and to isolate cDNA and genomic clones of other genes that have a high sequence similarity to these BSGs. Such probes 30 generally will comprise at least 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases.

For example, the coding region of BSG of the present invention may be isolated by screening using an 35 oligonucleotide probe synthesized from the known DNA sequence.

0986241-062101  
"Patent" 0986241-062101

A labeled oligonucleotide having a sequence complementary to that of a gene of the present invention is used to screen a library of human cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes with.

5           The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to human disease, as further discussed herein relating to polynucleotide assays, *inter alia*.

10           The polynucleotides may encode a polypeptide which is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a  
15 protein from precursor to a mature form, may facilitate/protein trafficking, may prolong or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in situ*, the additional amino acids may be processed away  
20 from the mature protein by cellular enzymes.

A precursor protein having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed, such inactive precursors generally are activated.  
25 Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

In sum, a polynucleotide of the present invention may encode a mature protein, a mature protein plus a leader  
30 sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences which are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which

09886241-062101

generally are removed during processing steps that produce active and mature forms of the polypeptide.

### ***Polypeptides***

The present invention further relates to BSG  
5 polypeptides, preferably polypeptides encoded by a  
polynucleotide of SEQ ID NO: 1, 2, 3, 4, or 5. An exemplary  
BSG polypeptide encoded by the BSG polynucleotide of SEQ ID  
NO:4 is depicted in SEQ ID NO:6. The invention also relates  
to fragments, analogs and derivatives of these polypeptides.  
10 The terms "fragment," "derivative" and "analog" when referring  
to the polypeptides of the present invention means a  
polypeptide which retains essentially the same biological  
function or activity as such polypeptides. Thus, an analog  
includes a proprotein which can be activated by cleavage of  
15 the proprotein portion to produce an active mature  
polypeptide.

The polypeptide of the present invention may be a  
recombinant polypeptide, a natural polypeptide or a synthetic  
polypeptide. In certain preferred embodiments it is a  
20 recombinant polypeptide.

The fragment, derivative or analog of a polypeptide of  
or the present invention may be (i) one in which one or more  
of the amino acid residues are substituted with a conserved  
or non-conserved amino acid residue (preferably a conserved  
25 amino acid residue) and such substituted amino acid residue  
may or may not be one encoded by the genetic code; (ii) one  
in which one or more of the amino acid residues includes a  
substituent group; (iii) one in which the mature polypeptide  
is fused with another compound, such as a compound to increase  
30 the half-life of the polypeptide (for example, polyethylene  
glycol); or (iv) one in which the additional amino acids are  
fused to the mature polypeptide, such as a leader or secretory  
sequence or a sequence which is employed for purification of  
the mature polypeptide or a proprotein sequence. Such

09886241-063101  
F0F290 T429860

fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

Among preferred variants are those that vary from a reference by conservative amino acid substitutions. Such  
5 substitutions are those that substitute a given amino acid in a polypeptide by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile; interchange of the hydroxyl residues  
10 Ser and Thr, exchange of the acidic residues Asp and Glu, substitution between the amide residues Asn and Gln, exchange of the basic residues Lys and Arg and replacements among the aromatic residues Phe, Tyr.

The polypeptides and polynucleotides of the present  
15 invention are preferably provided in an isolated form, and preferably are purified to homogeneity.

The polypeptides of the present invention include the polypeptides encoded by the polynucleotide of SEQ ID NO: 1, 2, 3, 4, or 5 (in particular the mature polypeptide) as well  
20 as polypeptides which have at least 75% similarity (preferably at least 75% identity), more preferably at least 90% similarity (more preferably at least 90% identity), still more preferably at least 95% similarity (still more preferably at least 95% identity), to a polypeptide encoded by SEQ ID NO:  
25 1, 2, 3, 4, or 5. Also included are portions of such polypeptides generally containing at least 30 amino acids and more preferably at least 50 amino acids.

As known in the art "similarity" between two polypeptides is determined by comparing the amino acid  
30 sequence and its conserved amino acid substitutes of one polypeptide sequence with that of a second polypeptide.

Fragments or portions of the polypeptides of the present invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, the  
35 fragments may be employed as intermediates for producing the

0986241-062101  
TOT290.T49866

full-length polypeptides. Fragments or portions of the polynucleotides of the present invention may be used to synthesize full-length polynucleotides of the present invention.

## 5 *Fragments*

Also among preferred embodiments of this aspect of the present invention are polypeptides comprising fragments of a polypeptide encoded by a polynucleotide of SEQ ID NO: 1, 2, 3, 4, or 5. In this regard a fragment is a polypeptide having  
10 an amino acid sequence that entirely is the same as part but not all of the amino acid sequence of the aforementioned BSG polypeptides and variants or derivatives thereof.

Such fragments may be "free-standing," i.e., not part of or fused to other amino acids or polypeptides, or they may  
15 be contained within a larger polypeptide of which they form a part or region. When contained within a larger polypeptide, the presently discussed fragments most preferably form a single continuous region. However, several fragments may be comprised within a single larger polypeptide. For instance,  
20 certain preferred embodiments relate to a fragment of a BSG polypeptide of the present comprised within a precursor polypeptide designed for expression in a host and having heterologous pre- and pro-polypeptide regions fused to the amino terminus of the BSG fragment and an additional region  
25 fused to the carboxyl terminus of the fragment. Therefore, fragments in one aspect of the meaning intended herein, refers to the portion or portions of a fusion polypeptide or fusion protein derived from a BSG polypeptide.

As representative examples of polypeptide fragments of  
30 the invention, there may be mentioned those which have from about 15 to about 139 amino acids. In this context "about" includes the particularly recited range and ranges larger or smaller by several, a few, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes. Highly preferred in this  
35 regard are the recited ranges plus or minus as many as 5 amino

09886241 "052101  
T0230 T42850

acids at either or at both extremes. Particularly highly preferred are the recited ranges plus or minus as many as 3 amino acids at either or at both the recited extremes. Especially preferred are ranges plus or minus 1 amino acid at either or at both extremes or the recited ranges with no additions or deletions. Most highly preferred of all in this regard are fragments from about 15 to about 45 amino acids.

Among especially preferred fragments of the invention are truncation mutants of the BSG polypeptides. Truncation mutants include BSG polypeptides having an amino acid sequence encoded by a polynucleotide of SEQ ID NO: 1, 2, 3, 4, or 5, or variants or derivatives thereof, except for deletion of a continuous series of residues (that is, a continuous region, part or portion) that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or, as in double truncation mutants, deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus. Fragments having the size ranges set out herein also are preferred embodiments of truncation fragments, which are especially preferred among fragments generally.

Also preferred in this aspect of the invention are fragments characterized by structural or functional attributes of the BSG polypeptides of the present invention. Preferred embodiments of the invention in this regard include fragments that comprise alpha-helix and alpha-helix forming regions ("alpha-regions"), beta-sheet and beta-sheet-forming regions ("beta-regions"), turn and turn-forming regions ("turn-regions"), coil and coil-forming regions ("coil-regions"), hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions and high antigenic index regions of the BSG polypeptides of the present invention. Regions of the aforementioned types are identified routinely by analysis of the amino acid sequences encoded by the polynucleotides of SEQ

0933241-062101  
TOT290" T4298360

ID NO: 1, 2, 3, 4, or 5. Preferred regions include Garnier-Robson alpha-regions, beta-regions, turn-regions and coil-regions, Chou-Fasman alpha-regions, beta-regions and turn-regions, Kyte-Doolittle hydrophilic regions and hydrophilic regions, Eisenberg alpha and beta amphipathic regions, Karplus-Schulz flexible regions, Emini surface-forming regions and Jameson-Wolf high antigenic index regions. Among highly preferred fragments in this regard are those that comprise regions of BSGs that combine several structural features, such as several of the features set out above. In this regard, the regions defined by selected residues of a BSG polypeptide which all are characterized by amino acid compositions highly characteristic of turn-regions, hydrophilic regions, flexible-regions, surface-forming regions, and high antigenic index-regions, are especially highly preferred regions. Such regions may be comprised within a larger polypeptide or may be by themselves a preferred fragment of the present invention, as discussed above. It will be appreciated that the term "about" as used in this paragraph has the meaning set out above regarding fragments in general.

Further preferred regions are those that mediate activities of BSG polypeptides. Most highly preferred in this regard are fragments that have a chemical, biological or other activity of a BSG polypeptide, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Highly preferred in this regard are fragments that contain regions that are homologs in sequence, or in position, or in both sequence and to active regions of related polypeptides, and which include breast specific-binding proteins. Among particularly preferred fragments in these regards are truncation mutants, as discussed above.

It will be appreciated that the invention also relates to polynucleotides encoding the aforementioned fragments, polynucleotides that hybridize to polynucleotides encoding the fragments, particularly those that hybridize under stringent

conditions, and polynucleotides such as PCR primers for amplifying polynucleotides that encode the fragments. In these regards, preferred polynucleotides are those that correspond to the preferred fragments, as discussed above.

## 5 ***Fusion Proteins***

In one embodiment of the present invention, the BSG polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of the present  
10 polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See also EP A 394,827; Traunecker, et al., Nature 331: 84-86 (1988)) Similarly, fusion to IgG-1, IgG-3, and albumin increases the halflife time *in vivo*. Nuclear localization signals fused to  
15 the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally,  
20 fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of these types of fusion proteins described above can be made in accordance with well known protocols.

For example, a BSG polypeptide can be fused to an IgG  
25 molecule via the following protocol. Briefly, the human Fc portion of the IgG molecule is PCR amplified using primers that span the 5' and 3' ends of the sequence. These primers also have convenient restriction enzyme sites that facilitate cloning into an expression vector, preferably a mammalian  
30 expression vector. For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. In this protocol, the 3' BamHI site must be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI thereby linearizing the vector,  
35 and a BSG polynucleotide of the present invention is ligated

09880241-062101  
TOT290-TH28560



into this BamHI site. It is preferred that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

If the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e. g., WO 96/34891.)

## 10 Diagnostic Assays

The present invention also relates to diagnostic assays and methods, both quantitative and qualitative for detecting, diagnosing, monitoring, staging and prognosticating cancers by comparing levels of BSG in a human patient with those of BSG in a normal human control. For purposes of the present invention, what is meant by BSG levels is, among other things, native protein expressed by a gene comprising the polynucleotide sequence of SEQ ID NO: 1, 2, 3, 4, or 5. An exemplary BSG protein encoded by the BSG polynucleotide of SEQ ID NO:4 is depicted in SEQ ID NO:6. By "BSG" it is also meant herein polynucleotides which, due to degeneracy in genetic coding, comprise variations in nucleotide sequence as compared to SEQ ID NO: 1, 2, 3, 4, or 5 but which still encode the same protein. The native protein being detected may be whole, a breakdown product, a complex of molecules or chemically modified. In the alternative, what is meant by BSG as used herein, means the native mRNA encoded by a polynucleotide sequence of SEQ ID NO: 1, 2, 3, 4, or 5, levels of the gene comprising the polynucleotide sequence of SEQ ID NO: 1, 2, 3, 4, or 5, or levels of a polynucleotide which is capable of hybridizing under stringent conditions to the antisense sequence of SEQ ID NO: 1, 2, 3, 4, or 5. Such levels are preferably determined in at least one of cells, tissues and/or bodily fluids, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in accordance

**0986471 = 00510**

with the invention for diagnosing overexpression of BSG protein compared to normal control bodily fluids, cells, or tissue samples may be used to diagnose the presence of breast cancer.

5 All the methods of the present invention may optionally include determining the levels of other cancer markers as well as BSG. Other cancer markers, in addition to BSG, useful in the present invention will depend on the cancer being tested and are known to those of skill in the art.

10           The present invention provides methods for diagnosing  
the presence of breast cancer by analyzing for changes in  
levels of BSG in cells, tissues or bodily fluids compared with  
levels of BSG in cells, tissues or bodily fluids of preferably  
the same type from a normal human control, wherein an increase  
15 in levels of BSG in the patient versus the normal human  
control is associated with the presence of breast cancer.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the patient being tested has cancer is one in which cells, tissues or bodily fluid levels of the cancer marker, such as BSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues or bodily fluid of a normal human control.

The present invention also provides a method of  
25 diagnosing metastatic breast cancer in a patient having breast  
cancer which has not yet metastasized for the onset of  
metastasis. In the method of the present invention, a human  
cancer patient suspected of having breast cancer which may  
have metastasized (but which was not previously known to have  
30 metastasized) is identified. This is accomplished by a  
variety of means known to those of skill in the art.

In the present invention, determining the presence of BSG levels in cells, tissues or bodily fluid, is particularly useful for discriminating between breast cancer which has not  
35 metastasized and breast cancer which has metastasized.

Existing techniques have difficulty discriminating between breast cancer which has metastasized and breast cancer which has not metastasized and proper treatment selection is often dependent upon such knowledge.

5 In the present invention, the cancer marker levels measured in such cells, tissues or bodily fluid is BSG, and are compared with levels of BSG in preferably the same cells, tissue or bodily fluid type of a normal human control. That is, if the cancer marker being observed is just BSG in serum,  
10 this level is preferably compared with the level of BSG in serum of a normal human control. An increase in the BSG in the patient versus the normal human control is associated with breast cancer which has metastasized.

Without limiting the instant invention, typically, for  
15 a quantitative diagnostic assay a positive result indicating the cancer in the patient being tested or monitored has metastasized is one in which cells, tissues or bodily fluid levels of the cancer marker, such as BSG, are at least two times higher, and most preferably are at least five times  
20 higher, than in preferably the same cells, tissues or bodily fluid of a normal patient.

Normal human control as used herein includes a human patient without cancer and/or non cancerous samples from the patient; in the methods for diagnosing or monitoring for  
25 metastasis, normal human control may preferably also include samples from a human patient that is determined by reliable methods to have breast cancer which has not metastasized.

### ***Staging***

The invention also provides a method of staging breast  
30 cancer in a human patient. The method comprises identifying a human patient having such cancer and analyzing cells, tissues or bodily fluid from such human patient for BSG. The BSG levels determined in the patient are then compared with levels of BSG in preferably the same cells, tissues or bodily  
35 fluid type of a normal human control, wherein an increase in

09336241-062401  
TOT290-1423360

BSG levels in the human patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of BSG (but still increased over true normal levels) is associated with a cancer which is regressing  
5 or in remission.

## Monitoring

Further provided is a method of monitoring breast cancer in a human patient having such cancer for the onset of metastasis. The method comprises identifying a human patient  
10 having such cancer that is not known to have metastasized; periodically analyzing cells, tissues or bodily fluid from such human patient for BSG; and comparing the BSG levels determined in the human patient with levels of BSG in preferably the same cells, tissues or bodily fluid type of a  
15 normal human control, wherein an increase in BSG levels in the human patient versus the normal human control is associated with a cancer which has metastasized. In this method, normal human control samples may also include prior patient samples.

Further provided by this invention is a method of  
20 monitoring the change in stage of breast cancer in a human  
patient having such cancer. The method comprises identifying  
a human patient having such cancer; periodically analyzing  
cells, tissues or bodily fluid from such human patient for  
BSG; and comparing the BSG levels determined in the human  
25 patient with levels of BSG in preferably the same cells,  
tissues or bodily fluid type of a normal human control,  
wherein an increase in BSG levels in the human patient versus  
the normal human control is associated with a cancer which is  
progressing in stage and a decrease in the levels of BSG is  
30 associated with a cancer which is regressing in stage or in  
remission. In this method, normal human control samples may  
also include prior patient samples.

Monitoring a patient for onset of metastasis is periodic and preferably done on a quarterly basis. However, this may

be done more or less frequently depending on the cancer, the particular patient, and the stage of the cancer.

***Prognostic Testing and Clinical Trial Monitoring***

The methods described herein can further be utilized as  
5 prognostic assays to identify subjects having or at risk of  
developing a disease or disorder associated with increased  
levels of BSG. The present invention provides a method in  
which a test sample is obtained from a human patient and BSG  
is detected. The presence of higher BSG levels as compared  
10 to normal human controls is diagnostic for the human patient  
being at risk for developing cancer, particularly breast  
cancer.

The effectiveness of therapeutic agents to decrease  
expression or activity of the BSGs of the invention can also  
15 be monitored by analyzing levels of expression of the BSGs in  
a human patient in clinical trials or in *in vitro* screening  
assays such as in human cells. In this way, the gene  
expression pattern can serve as a marker, indicative of the  
physiological response of the human patient, or cells as the  
20 case may be, to the agent being tested.

***Detection of genetic lesions or mutations***

The methods of the present invention can also be used  
to detect genetic lesions or mutations in BSG, thereby  
determining if a human with the genetic lesion is at risk for  
25 breast cancer or has breast cancer. Genetic lesions can be  
detected, for example, by ascertaining the existence of a  
deletion and/or addition and/or substitution of one or more  
nucleotides from the BSGs of this invention, a chromosomal  
rearrangement of BSG, aberrant modification of BSG (such as  
30 of the methylation pattern of the genomic DNA), the presence  
of a non-wild type splicing pattern of a mRNA transcript of  
BSG, allelic loss of BSG, and/or inappropriate post-  
translational modification of BSG protein. Methods to detect  
such lesions in the BSG of this invention are known to those  
35 of skill in the art.

09886241.069101

For example, in one embodiment, alterations in a gene corresponding to a BSG polynucleotide of the present invention are determined via isolation of RNA from entire families or individual patients presenting with a phenotype of interest (such as a disease) is be isolated. cDNA is then generated from these RNA samples using protocols known in the art. See, e.g. Sambrook et al. (MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)), which is illustrative of the many laboratory manuals that detail these techniques. The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO: 1, 2, 3, 4, or 5. PCR conditions typically consist of 35 cycles at 95°C for 30 seconds; 60-120 seconds at 52-58°C; and 60-120 seconds at 70°C, using buffer solutions described in Sidransky, D., et al., Science 252: 706 (1991). PCR products are sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase (Epicentre Technologies). The intron-exon borders of selected exons are also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations are then cloned and sequenced to validate the results of the direct sequencing. PCR products are cloned into T-tailed vectors as described in Holton, T. A. and Graham, M. W., Nucleic Acids Research, 19 : 1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

Genomic rearrangements can also be observed as a method of determining alterations in a gene corresponding to a polynucleotide. In this method, genomic clones are nick-translated with digoxigenin deoxy-uridine 5'triphosphate (Boehringer Mannheim), and FISH is performed as described in Johnson, C. et al., Methods Cell Biol. 35: 73-99 (1991). Hybridization with a labeled probe is carried out using a vast

excess of human DNA for specific hybridization to the corresponding genomic locus. Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C-and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters (Johnson et al., Genet. Anal. Tech. Appl., 8: 75 (1991)). Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System (Inovision Corporation, Durham, NC). Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated disease.

## Assay Techniques

Assay techniques that can be used to determine levels of gene expression (including protein levels), such as BSG of the present invention, in a sample derived from a patient are well known to those of skill in the art. Such assay methods include, without limitation, radioimmunoassays, reverse transcriptase PCR (RT-PCR) assays, immunohistochemistry assays, *in situ* hybridization assays, competitive-binding assays, Western Blot analyses, ELISA assays and proteomic approaches: two-dimensional gel electrophoresis (2D electrophoresis) and non-gel based approaches such as mass spectrometry or protein interaction profiling. Among these, ELISAs are frequently preferred to diagnose a gene's expressed protein in biological fluids.

30       An ELISA assay initially comprises preparing an antibody, if not readily available from a commercial source, specific to BSG, preferably a monoclonal antibody. In addition a reporter antibody generally is prepared which binds specifically to BSG. The reporter antibody is attached to a  
35 detectable reagent such as radioactive, fluorescent or

enzymatic reagent, for example horseradish peroxidase enzyme or alkaline phosphatase.

To carry out the ELISA, antibody specific to BSG is incubated on a solid support, e.g. a polystyrene dish, that binds the antibody. Any free protein binding sites on the dish are then covered by incubating with a non-specific protein such as bovine serum albumin. Next, the sample to be analyzed is incubated in the dish, during which time BSG binds to the specific antibody attached to the polystyrene dish. Unbound sample is washed out with buffer. A reporter antibody specifically directed to BSG and linked to a detectable reagent such as horseradish peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to BSG. Unattached reporter antibody is then washed out. Reagents for peroxidase activity, including a colorimetric substrate are then added to the dish. Immobilized peroxidase, linked to BSG antibodies, produces a colored reaction product. The amount of color developed in a given time period is proportional to the amount of BSG protein present in the sample. Quantitative results typically are obtained by reference to a standard curve.

A competition assay can also be employed wherein antibodies specific to BSG are attached to a solid support and labeled BSG and a sample derived from the host are passed over the solid support. The amount of label detected which is attached to the solid support can be correlated to a quantity of BSG in the sample.

Using all or a portion of a nucleic acid sequence of BSG of the present invention as a hybridization probe, nucleic acid methods can also be used to detect BSG mRNA as a marker for breast cancer. Polymerase chain reaction (PCR) and other nucleic acid methods, such as ligase chain reaction (LCR) and nucleic acid sequence based amplification (NASBA), can be used to detect malignant cells for diagnosis and monitoring of



various malignancies. For example, reverse-transcriptase PCR (RT-PCR) is a powerful technique which can be used to detect the presence of a specific mRNA population in a complex mixture of thousands of other mRNA species. In RT-PCR, an mRNA species is first reverse transcribed to complementary DNA (cDNA) with use of the enzyme reverse transcriptase; the cDNA is then amplified as in a standard PCR reaction. RT-PCR can thus reveal by amplification the presence of a single species of mRNA. Accordingly, if the mRNA is highly specific for the cell that produces it, RT-PCR can be used to identify the presence of a specific type of cell.

Hybridization to clones or oligonucleotides arrayed on a solid support (i.e. gridding) can be used to both detect the expression of and quantitate the level of expression of that gene. In this approach, a cDNA encoding the BSG gene is fixed to a substrate. The substrate may be of any suitable type including but not limited to glass, nitrocellulose, nylon or plastic. At least a portion of the DNA encoding the BSG gene is attached to the substrate and then incubated with the analyte, which may be RNA or a complementary DNA (cDNA) copy of the RNA, isolated from the tissue of interest. Hybridization between the substrate bound DNA and the analyte can be detected and quantitated by several means including but not limited to radioactive labeling or fluorescence labeling of the analyte or a secondary molecule designed to detect the hybrid. Quantitation of the level of gene expression can be done by comparison of the intensity of the signal from the analyte compared with that determined from known standards. The standards can be obtained by *in vitro* transcription of the target gene, quantitating the yield, and then using that material to generate a standard curve.

Of the proteomic approaches, 2D electrophoresis is a technique well known to those in the art. Isolation of individual proteins from a sample such as serum is accomplished using sequential separation of proteins by

09886241-062101

different characteristics usually on polyacrylamide gels. First, proteins are separated by size using an electric current. The current acts uniformly on all proteins, so smaller proteins move farther on the gel than larger proteins.

5 The second dimension applies a current perpendicular to the first and separates proteins not on the basis of size but on the specific electric charge carried by each protein. Since no two proteins with different sequences are identical on the basis of both size and charge, the result of a 2D separation  
10 is a square gel in which each protein occupies a unique spot. Analysis of the spots with chemical or antibody probes, or subsequent protein microsequencing can reveal the relative abundance of a given protein and the identity of the proteins in the sample.

15 The above tests can be carried out on samples derived from a variety of cells, bodily fluids and/or tissue extracts such as homogenates or solubilized tissue obtained from a patient. Tissue extracts are obtained routinely from tissue biopsy and autopsy material. Bodily fluids useful in the  
20 present invention include blood, urine, saliva or any other bodily secretion or derivative thereof. By blood it is meant to include whole blood, plasma, serum or any derivative of blood.

#### ***In Vivo Targeting of BSG/Breast Cancer Therapy***

25 Identification of this BSG is also useful in the rational design of new therapeutics for imaging and treating cancers, and in particular breast cancer. For example, in one embodiment, antibodies which specifically bind to BSG can be raised and used *in vivo* in patients suspected of suffering  
30 from breast cancer. Antibodies which specifically bind BSG can be injected into a patient suspected of having breast cancer for diagnostic and/or therapeutic purposes. Thus, another aspect of the present invention provides for a method for preventing the onset and treatment of breast cancer in a  
35 human patient in need of such treatment by administering to

0986241-052101

[illegible]

Antibodies which can be used in *in vivo* methods include polyclonal, monoclonal and omniclonal antibodies and antibodies prepared via molecular biology techniques. Antibody fragments and aptamers and single-stranded  
5 oligonucleotides such as those deriyed from an *in vitro* evolution protocol referred to as SELEX and well known to those skilled in the art can also be used.

#### **Screening Assays**

The present invention also provides methods for  
10 identifying modulators which bind to BSG protein or have a modulatory effect on the expression or activity of BSG protein. Modulators which decrease the expression or activity of BSG protein are believed to be useful in treating breast cancer. Such screening assays are known to those of skill in  
15 the art and include, without limitation, cell-based assays and cell free assays.

Small molecules predicted via computer imaging to specifically bind to regions of BSG can also be designed, synthesized and tested for use in the imaging and treatment  
20 of breast cancer. Further, libraries of molecules can be screened for potential anticancer agents by assessing the ability of the molecule to bind to the BSGs identified herein. Molecules identified in the library as being capable of binding to BSG are key candidates for further evaluation for  
25 use in the treatment of breast cancer. In a preferred embodiment, these molecules will downregulate expression and/or activity of BSG in cells.

#### **Adoptive Immunotherapy and Vaccines**

Adoptive immunotherapy of cancer refers to a therapeutic  
30 approach in which immune cells with an antitumor reactivity are administered to a tumor-bearing host, with the aim that the cells mediate either directly or indirectly, the regression of an established tumor. Transfusion of lymphocytes, particularly T lymphocytes, falls into this  
35 category and investigators at the National Cancer Institute

09886241.062101

(NCI) have used autologous reinfusion of peripheral blood lymphocytes or tumor-infiltrating lymphocytes (TIL), T cell cultures from biopsies of subcutaneous lymph nodules, to treat several human cancers (Rosenberg, S. A., U.S. Patent No. 5 4,690,914, issued Sep. 1, 1987; Rosenberg, S. A., et al., 1988, N. England J. Med. 319:1676-1680).

09886241-062101  
The present invention relates to compositions and methods of adoptive immunotherapy for the prevention and/or treatment of primary and metastatic breast cancer in humans 10 using macrophages sensitized to the antigenic BSG molecules, with or without non-covalent complexes of heat shock protein (hsp). Antigenicity or immunogenicity of the BSG is readily confirmed by the ability of the BSG protein or a fragment thereof to raise antibodies or educate naive effector cells, 15 which in turn lyse target cells expressing the antigen (or epitope).

Cancer cells are, by definition, abnormal and contain proteins which should be recognized by the immune system as foreign since they are not present in normal tissues. However, 20 the immune system often seems to ignore this abnormality and fails to attack tumors. The foreign BSG proteins that are produced by the cancer cells can be used to reveal their presence. The BSG is broken into short fragments, called tumor antigens, which are displayed on the surface of the 25 cell. These tumor antigens are held or presented on the cell surface by molecules called MHC, of which there are two types: class I and II. Tumor antigens in association with MHC class I molecules are recognized by cytotoxic T cells while antigen-MHC class II complexes are recognized by a second subset of 30 T cells called helper cells. These cells secrete cytokines which slow or stop tumor growth and help another type of white blood cell, B cells, to make antibodies against the tumor cells.

In adoptive immunotherapy, T cells or other antigen 35 presenting cells (APCs) are stimulated outside the body (ex

vivo), using the tumor specific BSG antigen. The stimulated cells are then reinfused into the patient where they attack the cancerous cells. Research has shown that using both cytotoxic and helper T cells is far more effective than using  
5 either subset alone. Additionally, the BSG antigen may be complexed with heat shock proteins to stimulate the APCs as described in U.S. Patent No. 5,985,270.

The APCs can be selected from among those antigen presenting cells known in the art including, but not limited  
10 to, macrophages, dendritic cells, B lymphocytes, and a combination thereof, and are preferably macrophages. In a preferred use, wherein cells are autologous to the individual, autologous immune cells such as lymphocytes, macrophages or other APCs are used to circumvent the issue of whom to select  
15 as the donor of the immune cells for adoptive transfer. Another problem circumvented by use of autologous immune cells is graft versus host disease which can be fatal if unsuccessfully treated.

In adoptive immunotherapy with gene therapy, DNA of the  
20 BSG can be introduced into effector cells similarly as in conventional gene therapy. This can enhance the cytotoxicity of the effector cells to tumor cells as they have been manipulated to produce the antigenic protein resulting in improvement of the adoptive immunotherapy.

BSG antigens of this invention are also useful as  
25 components of breast cancer vaccines. The vaccine comprises an immunogenically stimulatory amount of a BSG antigen. Immunogenically stimulatory amount refers to that amount of antigen that is able to invoke the desired immune response in  
30 the recipient for the amelioration, or treatment of breast cancer. Effective amounts may be determined empirically by standard procedures well known to those skilled in the art.

The BSG antigen may be provided in any one of a number of vaccine formulations which are designed to induce the  
35 desired type of immune response, e.g., antibody and/or cell

09886241.062101

mediated. Such formulations are known in the art and include, but are not limited to, formulations such as those described in U.S. Patent 5,585,103. Vaccine formulations of the present invention used to stimulate immune responses can also include  
5 pharmaceutically acceptable adjuvants.

***Vectors, host cells, expression***

The present invention also relates to vectors which include polynucleotides of the present invention, host cells which are genetically engineered with vectors of the invention  
10 and the production of polypeptides of the invention by recombinant techniques.

Host cells can be genetically engineered to incorporate BSG polynucleotides and express BSG polypeptides of the present invention. For instance, BSG polynucleotides may be  
15 introduced into host cells using well known techniques of infection, transduction, transfection, transvection and transformation. The BSG polynucleotides may be introduced alone or with other polynucleotides. Such other polynucleotides may be introduced independently, co-introduced  
20 or introduced joined to the BSG polynucleotides of the invention.

For example, BSG polynucleotides of the invention may be transfected into host cells with another, separate, polynucleotide encoding a selectable marker, using standard  
25 techniques for co-transfection and selection in, for instance, mammalian cells. In this case, the polynucleotides generally will be stably incorporated into the host cell genome.

Alternatively, the BSG polynucleotide may be joined to a vector containing a selectable marker for propagation in a  
30 host. The vector construct may be introduced into host cells by the aforementioned techniques. Generally, a plasmid vector is introduced as DNA in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. Electroporation also may be used to introduce BSG  
35 polynucleotides into a host. If the vector is a virus, it may

09886241-052101  
TOTAL: 290

be packaged *in vitro* or introduced into a packaging cell and the packaged virus may be transduced into cells. A wide variety of well known techniques conducted routinely by those of skill in the art are suitable for making BSG  
5 polynucleotides and for introducing BSG polynucleotides into cells in accordance with this aspect of the invention. Such techniques are reviewed at length in reference texts such as Sambrook et al., previously cited herein.

Vectors which may be used in the present invention  
10 include, for example, plasmid vectors, single- or double-stranded phage vectors, and single- or double-stranded RNA or DNA viral vectors. Such vectors may be introduced into cells as polynucleotides, preferably DNA, by well known techniques for introducing DNA and RNA into cells. The vectors, in the  
15 case of phage and viral vectors, also may be and preferably are introduced into cells as packaged or encapsidated virus by well known techniques for infection and transduction. Viral vectors may be replication competent or replication defective. In the latter case viral propagation generally  
20 will occur only in complementing host cells.

Preferred vectors for expression of polynucleotides and polypeptides of the present invention include, but are not limited to, vectors comprising cis-acting control regions effective for expression in a host operatively linked to the  
25 polynucleotide to be expressed. Appropriate trans-acting factors either are supplied by the host, supplied by a complementing vector or supplied by the vector itself upon introduction into the host.

In certain preferred embodiments in this regard, the  
30 vectors provide for specific expression. Such specific expression may be inducible expression or expression only in certain types of cells or both inducible and cell-specific. Particularly preferred among inducible vectors are vectors that can be induced to express by environmental factors that  
35 are easy to manipulate, such as temperature and nutrient

00886241 092101  
101230 142360



additives. A variety of vectors suitable to this aspect of the invention, including constitutive and inducible expression vectors for use in prokaryotic and eukaryotic hosts, are well known and employed routinely by those of skill in the art.

5 The engineered host cells can be cultured in conventional nutrient media which may be modified as appropriate for, *inter alia*, activating promoters, selecting transformants or amplifying genes. Culture conditions such as temperature, pH and the like, previously used with the host  
10 cell selected for expression, generally will be suitable for expression of BSG polypeptides of the present invention.

A great variety of expression vectors can be used to express BSG polypeptides of the invention. Such vectors include chromosomal, episomal and virus-derived vectors.  
15 Vectors may be derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and from  
20 combinations thereof such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. All may be used for expression in accordance with this aspect of the present invention. Generally, any vector suitable to maintain, propagate or express polynucleotides to express a  
25 polypeptide in a host may be used for expression in this regard.

The appropriate DNA sequence may be inserted into the vector by any of a variety of well-known and routine techniques. In general, a DNA sequence for expression is  
30 joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more restriction endonucleases and then joining the restriction fragments together using T4 DNA ligase. Procedures for restriction and ligation that can be used to this end are well known and  
35 routine to those of skill. Suitable procedures in this

0938241-06101  
FOR "99" T29860

regard, and for constructing expression vectors using alternative techniques, which also are well known and routine to those skill, are set forth in great detail in Sambrook et al. cited elsewhere herein.

5       The DNA sequence in the expression vector is operatively linked to appropriate expression control sequence(s), including, for instance, a promoter to direct mRNA transcription. Representative promoters include the phage  
10   lambda PL promoter, the *E. coli* lac, trp and tac promoters, the SV40 early and late promoters, and promoters of retroviral LTRs, to name just a few of the well-known promoters. It will be understood that numerous promoters not mentioned are also  
15   suitable for use in this aspect of the invention and are well known and readily may be employed by those of skill in the manner illustrated by the discussion and the examples herein.

      In general, expression constructs will contain sites for transcription initiation and termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts expressed by the  
20   constructs will include a translation initiating AUG at the beginning and a termination codon appropriately positioned at the end of the polypeptide to be translated.

      In addition, the constructs may contain control regions that regulate as well as engender expression. Generally, in  
25   accordance with many commonly practiced procedures, such regions will operate by controlling transcription, such as repressor binding sites and enhancers, among others.

      Vectors for propagation and expression generally will include selectable markers. Such markers also may be suitable  
30   for amplification or the vectors may contain additional markers for this purpose. In this regard, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells. Preferred markers include dihydrofolate reductase  
35   or neomycin resistance for eukaryotic cell culture, and

09886241-062101

tetracycline or ampicillin resistance genes for culturing in *E. coli* and other bacteria.

The vector containing the appropriate DNA sequence as described elsewhere herein, as well as an appropriate promoter, and other appropriate control sequences, may be introduced into an appropriate host using a variety of well known techniques suitable to expression therein of a desired polypeptide. Representative examples of appropriate hosts include bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS and Bowes melanoma cells; and plant cells. Hosts for a great variety of expression constructs are well known, and those of skill will be enabled by the present disclosure readily to select a host for expressing a BSG polypeptide in accordance with this aspect of the present invention.

More particularly, the present invention also includes recombinant constructs, such as expression constructs, comprising one or more of the sequences described above. The constructs comprise a vector, such as a plasmid or viral vector, into which such BSG sequence of the invention has been inserted. The sequence may be inserted in a forward or reverse orientation. In certain preferred embodiments in this regard, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and there are many commercially available vectors suitable for use in the present invention.

The following vectors, which are commercially available, are provided by way of example. Among vectors preferred for use in bacteria are pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and

ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are PWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. These  
5 vectors are listed solely by way of illustration of the many commercially available and well known vectors that are available to those of skill in the art for use in accordance with this aspect of the present invention. It will be appreciated by those of skill in the art upon reading this  
10 disclosure that any other plasmid or vector suitable for introduction, maintenance, propagation and/or expression of a BSG polynucleotide or polypeptide of the invention in a host may be used in this aspect of the invention.

Promoter regions can be selected from any desired gene  
15 using vectors that contain a reporter transcription unit lacking a promoter region, such as a chloramphenicol acetyl transferase ("cat") transcription unit, downstream of a restriction site or sites for introducing a candidate promoter fragment; i.e., a fragment that may contain a promoter. As  
20 is well known, introduction into the vector of a promoter-containing fragment at the restriction site upstream of the cat gene engenders production of CAT activity detectable by standard CAT assays. Vectors suitable to this end are well known and readily available. Two such vectors are pKK232-8  
25 and pCM7. Thus, promoters for expression of BSG polynucleotides of the present invention include, not only well known and readily available promoters, but also promoters that readily may be obtained by the foregoing technique, using a reporter gene.

30 Among known bacterial promoters suitable for expression of polynucleotides and polypeptides in accordance with the present invention are the *E. coli* lacI and lacZ promoters, the T3 and T7 promoters, the gpt promoter, the lambda PR, PL promoters and the trp promoter. Among known eukaryotic  
35 promoters suitable in this regard are the CMV immediate early

09886241-062101

promoter, the HSV thymidine kinase promoter, the early and late SV40 promoters, the promoters of retroviral LTRs, such as those of the Rous sarcoma virus ("RSV"), and metallothionein promoters, such as the mouse metallothionein-I  
5 promoter.

Selection of appropriate vectors and promoters for expression in a host cell is a well known procedure and the requisite techniques for expression vector construction, introduction of the vector into the host and expression in the  
10 host are routine skills in the art.

The present invention also relates to host cells containing the above-described constructs. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell. Alternatively,  
15 the host cell can be a prokaryotic cell, such as a bacterial cell.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods.  
20 Such methods are described in many standard laboratory manuals, such as Davis et al. BASIC METHODS IN MOLECULAR BIOLOGY, (1986).

Constructs in host cells can be used in a conventional  
25 manner to produce the gene product encoded by the recombinant sequence. Alternatively, BSG polypeptides of the invention can be synthetically produced by conventional peptide synthesizers.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of  
30 appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and

09866241-062401

eukaryotic hosts are described by Sambrook et al. cited elsewhere herein.

Generally, recombinant expression vectors will include origins of replication, a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence, and a selectable marker to permit isolation of vector containing cells after exposure to the vector. Among suitable promoters are those derived from the genes that encode glycolytic enzymes such as 3-phosphoglycerate kinase ("PGK"),  $\alpha$ -factor, acid phosphatase, and heat shock proteins, among others. Selectable markers include the ampicillin resistance gene of *E. coli* and the *trp1* gene of *S. cerevisiae*.

Transcription of DNA encoding the BSG polypeptides of the present invention by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 base pairs (bp) that act to increase transcriptional activity of a promoter in a given host cell-type. Examples of enhancers include the SV40 enhancer, which is located on the late side of the replication origin at bp 100 to 270, the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

A polynucleotide of the present invention, encoding a heterologous structural sequence of a BSG polypeptide of the present invention, generally will be inserted into the vector using standard techniques so that it is operably linked to the promoter for expression. The polynucleotide will be positioned so that the transcription start site is located appropriately 5' to a ribosome binding site. The ribosome binding site will be 5' to the AUG that initiates translation of the polypeptide to be expressed. Generally, there will be no other open reading frames that begin with an initiation codon, usually AUG, lying between the ribosome binding site

09886241.062101  
TOT290" T429860

and the initiating AUG. Also, generally, there will be a translation stop codon at the end of the polypeptide and there will be a polyadenylation signal and a transcription termination signal appropriately disposed at the 3' end of the  
5 transcribed region.

Appropriate secretion signals may be incorporated into the expressed polypeptide for secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment. The  
10 signals may be endogenous to the polypeptide or they may be heterologous signals.

The polypeptide may be expressed in a modified form, such as a fusion protein, and may include not only secretion signals but also additional heterologous functional regions.  
15 Thus, for instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence in the host cell during purification or during subsequent handling and storage. A region also may be added  
20 to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to polypeptides to engender secretion or excretion, to improve stability and to facilitate purification, among others, are familiar and routine  
25 techniques in the art.

Suitable prokaryotic hosts for propagation, maintenance or expression of BSG polynucleotides and polypeptides in accordance with the invention include *Escherichia coli*, *Bacillus subtilis* and *Salmonella typhimurium*. Various species  
30 of *Pseudomonas*, *Streptomyces*, and *Staphylococcus* are suitable hosts in this regard. Many other hosts also known to those of skill may also be employed in this regard.

As a representative, but non-limiting example, useful expression vectors for bacterial use can comprise a selectable  
35 marker and bacterial origin of replication derived from

09336241-062101  
101230-143360

commercially available plasmids comprising genetic elements of the well known cloning vector pBR322. Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, Wis., USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, where the selected promoter is inducible it is induced by appropriate means (e.g., temperature shift or exposure to chemical inducer) and cells are cultured for an additional period. Cells typically then are harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification. Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, such methods are well known to those skilled in the art.

Various mammalian cell culture systems can be employed for expression, as well. An exemplary mammalian expression system is the COS-7 line of monkey kidney fibroblasts described in Gluzman et al., Cell 23: 175 (1981). Other mammalian cell lines capable of expressing a compatible vector include for example, the C127, 3T3, CHO, HeLa, human kidney 293 and BHK cell lines. Mammalian expression vectors comprise an origin of replication, a suitable promoter and enhancer, and any ribosome binding sites, polyadenylation sites, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking non-transcribed sequences that are necessary for expression. In certain preferred embodiments in this regard DNA sequences derived from the SV40 splice sites, and the SV40 polyadenylation sites are used for required non-transcribed genetic elements of these types.



BSG polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

BSG polypeptides of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the BSG polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, BSG polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes.

BSG polynucleotides and polypeptides may be used in accordance with the present invention for a variety of applications, particularly those that make use of the chemical and biological properties of the BSGs. Additional applications relate to diagnosis and to treatment of disorders of cells, tissues and organisms. These aspects of the invention are illustrated further by the following discussion.

### **Polynucleotide assays**

As discussed in some detail *supra*, this invention is also related to the use of BSG polynucleotides to detect complementary polynucleotides such as, for example, as a diagnostic reagent. Detection of a mutated form of BSG associated with a dysfunction will provide a diagnostic tool

that can add to or define a diagnosis of a disease or susceptibility to a disease which results from under-expression, over-expression or altered expression of a BSG, such as, for example, a susceptibility to inherited breast  
5 cancer.

Individuals carrying mutations in a human BSG gene may be detected at the DNA level by a variety of techniques. Nucleic acids for diagnosis may be obtained from a patient's cells, such as from blood, urine, saliva, tissue biopsy and  
10 autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically using PCR prior to analysis (Saiki et al., Nature, 324: 163-166 (1986)). RNA or cDNA may also be used in a similar manner. As an example, PCR primers complementary to a BSG polynucleotide of SEQ ID  
15 NO: 1, 2, 3, 4, or 5 can be used to identify and analyze BSG expression and mutations. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to  
20 radiolabeled BSG RNA or alternatively, radiolabeled BSG antisense DNA sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures.

Sequence differences between a reference gene and genes  
25 having mutations also may be revealed by direct DNA sequencing. In addition, cloned DNA segments may be employed as probes to detect specific DNA segments. The sensitivity of such methods can be greatly enhanced by appropriate use of PCR or another amplification method. For example, a  
30 sequencing primer is used with double-stranded PCR product or a single-stranded template molecule generated by a modified PCR. The sequence determination is performed by conventional procedures with radiolabeled nucleotide or by automatic sequencing procedures with fluorescent-tags.

09885241-062101

Genetic testing based on DNA sequence differences may be achieved by detection of alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents. Small sequence deletions and insertions can be visualized by high resolution gel electrophoresis. DNA fragments of different sequences may be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures (see, e.g., Myers et al., Science, 230: 1242 (1985)).

Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (e.g., Cotton et al., Proc. Natl. Acad. Sci., USA, 85: 4397-4401 (1985)).

Thus, the detection of a specific DNA sequence may be achieved by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing or the use of restriction enzymes, (e.g., restriction fragment length polymorphisms ("RFLP") and Southern blotting of genomic DNA. In addition to more conventional gel-electrophoresis and DNA sequencing, mutations also can be detected by *in situ* analysis.

#### **Chromosome assays**

The BSG sequences of the present invention are also valuable for chromosome identification. There is a need for identifying particular sites on the chromosome and few chromosome marking reagents based on actual sequence data (repeat polymorphisms) are presently available for marking chromosomal location. Each BSG sequence of the present invention is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Thus, the BSGs can be used in the mapping of DNAs to chromosomes, an important first step in correlating sequences with genes associated with disease.

In certain preferred embodiments in this regard, the cDNA herein disclosed is used to clone genomic DNA of a BSG of the present invention. This can be accomplished using a variety of well known techniques and libraries, which  
5 generally are available commercially. The genomic DNA is used for *in situ* chromosome mapping using well known techniques for this purpose.

In some cases, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA.  
10 Computer analysis of the 3' untranslated region of the gene is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human  
15 chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified fragment.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide  
20 primers, sublocalization can be achieved with panels of fragments from specific chromosomes or pools of large genomic clones in an analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include *in situ* hybridization, prescreening with labeled flow-sorted  
25 chromosomes and preselection by hybridization to construct chromosome specific-cDNA libraries.

Fluorescence *in situ* hybridization ("FISH") of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique  
30 can be used with cDNA as short as 50 or 60 bp. This technique is described by Verma et al. (HUMAN CHROMOSOMES: A MANUAL OF BASIC TECHNIQUES, Pergamon Press, New York (1988)).

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the  
35 chromosome can be correlated with genetic map data. Such data

09886241.062101

are found, for example, in V. McKusick, MENDELIAN INHERITANCE IN MAN, available on line through Johns Hopkins University, Welch Medical Library. The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

#### ***Polypeptide assays***

As described in some detail *supra*, the present invention also relates to diagnostic assays such as quantitative and diagnostic assays for detecting levels of BSG polypeptide in cells and tissues, and biological fluids such as blood and urine, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in accordance with the present invention for detecting over-expression or under-expression of a BSG polypeptide compared to normal control tissue samples may be used to detect the presence of neoplasia. Assay techniques that can be used to determine levels of a protein, such as a BSG polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays. Among these ELISAs frequently are preferred.

For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10  $\mu\text{g/ml}$ . The antibodies are either monoclonal or polyclonal and are produced by methods as described herein. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced. The coated wells are then incubated for > 2 hours at room temperature with a sample containing the BSG polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded polypeptide. Next, 50  $\mu\text{l}$  of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate. 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution (75 $\mu\text{l}$ ) is then added to each well and the plate is incubated 1 hour at room temperature. The reaction is measured by a microtiter plate reader. A standard curve is prepared using serial dilutions of a control sample, and polypeptide concentration is plotted on the X-axis (log scale) while fluorescence or absorbance is plotted on the Y-axis (linear scale). The concentration of the BSG polypeptide in the sample is interpolated using the standard curve.

#### **Antibodies**

As discussed in some detail *supra*, BSG polypeptides, their fragments or other derivatives, or analogs thereof, or cells expressing them can be used as an immunogen to produce antibodies thereto. These antibodies can be polyclonal or monoclonal antibodies. The present invention also includes chimeric, single chain, and humanized antibodies, as well as Fab fragments, or the product of an Fab expression library.

Various procedures known in the art may be used for the production of such antibodies and fragments.

A variety of methods for antibody production are set forth in Current Protocols, Chapter 2.

5 For example, cells expressing a BSG polypeptide of the present invention can be administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of the secreted protein is prepared and purified to render it substantially free of  
10 natural contaminants. This preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity. The antibody obtained will bind with the BSG polypeptide itself. In this manner, even a sequence encoding only a fragment of the BSG polypeptide can  
15 be used to generate antibodies binding the whole native polypeptide. Such antibodies can then be used to isolate the BSG polypeptide from tissue expressing that BSG polypeptide.

Alternatively, monoclonal antibodies can be prepared. Examples of techniques for production of monoclonal antibodies  
20 include, but are not limited to, the hybridoma technique (Kohler, G. and Milstein, C., Nature 256: 495-497 (1975), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., Immunology Today 4: 72 (1983) and (Cole et al., pg. 77-96 in MONOCLONAL ANTIBODIES AND CANCER THERAPY, Alan R.  
25 Liss, Inc. (1985). The EBV-hybridoma technique is useful in production of human monoclonal antibodies.

Hybridoma technologies have also been described by Khler et al. (Eur. J. Immunol. 6: 511 (1976)) Khler et al. (Eur. J. Immunol. 6: 292 (1976)) and Hammerling et al. (in:  
30 Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N. Y., pp. 563-681 (1981)). In general, such procedures involve immunizing an animal (preferably a mouse) with BSG polypeptide or, more preferably, with a secreted BSG polypeptide-expressing cell. Such cells may be cultured in any suitable  
35 tissue culture medium; however, it is preferable to culture

09886241-062101

cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin. The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP20), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80: 225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide.

Alternatively, additional antibodies capable of binding to the polypeptide can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the protein-specific antibody can be blocked by the polypeptide. Such antibodies comprise anti-idiotypic antibodies to the protein specific antibody and can be used to immunize an animal to induce formation of further protein-specific antibodies.

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can also be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention. Also, transgenic mice, as well as other nonhuman transgenic animals, may be used to express



humanized antibodies to immunogenic polypeptide products of this invention.

It will be appreciated that Fab, F(ab')<sub>2</sub> and other fragments of the antibodies of the present invention may also be used according to the methods disclosed herein. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')<sub>2</sub> fragments). Alternatively, secreted protein-binding fragments can be produced through the application of recombinant DNA technology or through synthetic chemistry.

For *in vivo* use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in the art (See, for review, Morrison, Science 229: 1202 (1985); Oi et al., BioTechniques 4: 214 (1986); Cabilly et al., U. S. Patent 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312: 643 (1984); Neuberger et al., Nature 314: 268 (1985).)

The above-described antibodies may be employed to isolate or to identify clones expressing BSG polypeptides or purify BSG polypeptides of the present invention by attachment of the antibody to a solid support for isolation and/or purification by affinity chromatography. As discussed in more detail *supra*, antibodies specific against a BSG may also be used to image tumors, particularly cancer of the breast, in patients suffering from cancer. Such antibodies may also be used therapeutically to target tumors expressing a BSG.

#### ***BSG binding molecules and assays***

This invention also provides a method for identification of molecules, such as receptor molecules, that bind BSGs.

Genes encoding proteins that bind BSGs, such as receptor proteins, can be identified by numerous methods known to those of skill in the art. Examples include, but are not limited to, ligand panning and FACS sorting. Such methods are described in many laboratory manuals such as, for instance, Coligan et al., Current Protocols in Immunology 1(2): Chapter 5 (1991).

Expression cloning may also be employed for this purpose. To this end, polyadenylated RNA is prepared from a cell responsive to a BSG of the present invention. A cDNA library is created from this RNA and the library is divided into pools. The pools are then transfected individually into cells that are not responsive to a BSG of the present invention. The transfected cells then are exposed to labeled BSG. BSG polypeptides can be labeled by a variety of well-known techniques including, but not limited to, standard methods of radio-iodination or inclusion of a recognition site for a site-specific protein kinase. Following exposure, the cells are fixed and binding of labeled BSG is determined. These procedures conveniently are carried out on glass slides. Pools containing labeled BSG are identified as containing cDNA that produced BSG-binding cells. Sub-pools are then prepared from these positives, transfected into host cells and screened as described above. Using an iterative sub-pooling and re-screening process, one or more single clones that encode the putative binding molecule, such as a receptor molecule, can be isolated.

Alternatively a labeled ligand can be photoaffinity linked to a cell extract, such as a membrane or a membrane  
30 extract, prepared from cells that express a molecule that it binds, such as a receptor molecule. Cross-linked material is resolved by polyacrylamide gel electrophoresis ("PAGE") and exposed to X-ray film. The labeled complex containing the ligand-receptor can be excised, resolved into peptide  
35 fragments, and subjected to protein microsequencing. The

amino acid sequence obtained from microsequencing can be used to design unique or degenerate oligonucleotide probes to screen cDNA libraries to identify genes encoding the putative receptor molecule.

- 5 Polypeptides of the invention also can be used to assess BSG binding capacity of BSG binding molecules, such as receptor molecules, in cells or in cell-free preparations.

***Agonists and antagonists - assays and molecules***

- The invention also provides a method of screening  
10 compounds to identify those which enhance or block the action of a BSG on cells. By "compound", as used herein, it is meant to be inclusive of small organic molecules, peptides, polypeptides and antibodies as well as any other candidate molecules which have the potential to enhance or agonize or  
15 block or antagonize the action of BSG on cells. As used herein, an agonist is a compound which increases the natural biological functions of a BSG or which functions in a manner similar to a BSG, while an antagonist, as used herein, is a compound which decreases or eliminates such functions.  
20 Various known methods for screening for agonists and/or antagonists can be adapted for use in identifying BSG agonist or antagonists.

- For example, a cellular compartment, such as a membrane or a preparation thereof, such as a membrane-preparation, may  
25 be prepared from a cell that expresses a molecule that binds a BSG, such as a molecule of a signaling or regulatory pathway modulated by BSG. The preparation is incubated with labeled BSG in the absence or the presence of a compound which may be a BSG agonist or antagonist. The ability of the compound to  
30 bind the binding molecule is reflected in decreased binding of the labeled ligand. Compounds which bind gratuitously, i.e., without inducing the effects of a BSG upon binding to the BSG binding molecule are most likely to be good antagonists. Compounds that bind well and elicit effects that  
35 are the same as or closely related to BSG are agonists. BSG-

like effects of potential agonists and antagonists may be measured, for instance, by determining activity of a second messenger system following interaction of the candidate molecule with a cell or appropriate cell preparation, and comparing the effect with that of BSG or molecules that elicit the same effects as BSG. Second messenger systems that may be useful in this regard include, but are not limited to, AMP guanylate cyclase, ion channel or phosphoinositide hydrolysis second messenger systems.

Another example of an assay for BSG antagonists is a competitive assay that combines BSG and a potential antagonist with membrane-bound BSG receptor molecules or recombinant BSG receptor molecules under appropriate conditions for a competitive inhibition assay. BSG can be labeled, such as by radioactivity, such that the number of BSG molecules bound to a receptor molecule can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a BSG polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a receptor molecule, without inducing BSG-induced activities, thereby preventing the action of BSG by excluding BSG from binding.

Potential antagonists include small molecules which bind to and occupy the binding site of the BSG polypeptide thereby preventing binding to cellular binding molecules, such as receptor molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules.

Other potential antagonists include antisense molecules.  
35 Antisense technology can be used to control gene expression

098641-06101

through antisense DNA or RNA or through triple-helix formation. Antisense techniques are discussed, for example, in Okano, J. Neurochem. 56: 560 (1991); OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION, CRC Press, Boca Raton, Fla. (1988). Triple helix formation is discussed in, for instance Lee et al., Nucleic Acids Research 6: 3073 (1979); Cooney et al., Science 241: 456 (1988); and Dervan et al., Science 251: 1360 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA. For example, the 5' coding portion of a polynucleotide that encodes a mature BSG polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of a BSG polypeptide. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into a BSG polypeptide. The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of a BSG.

#### **Compositions**

The present invention also relates to compositions comprising a BSG polynucleotide or a BSG polypeptide or an agonist or antagonist thereof.

For example, a BSG polynucleotide, polypeptide or an agonist or antagonist thereof of the present invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water,

glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration.

Compositions of the present invention will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the polypeptide or other compound alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of secreted polypeptide administered parenterally per dose will be in the range of about 1,  $\mu\text{g/kg/day}$  to 10  $\text{mg/kg/day}$  of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01  $\text{mg/kg/day}$ , and most preferably for humans between about 0.01 and 1  $\text{mg/kg/day}$  for the hormone. If given continuously, the polypeptide or other compound is typically administered at a dose rate of about 1  $\mu\text{g/kg/hour}$  to about 50  $\text{mg/kg/hour}$ , either by 1-4 injections per day or by continuous subcutaneous infusion, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Pharmaceutical compositions containing the secreted protein of the invention are administered orally, rectally, parenterally, intracistemally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

5       The polypeptide or other compound is also suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include semipermeable polymer matrices in the form of shaped articles, e. g., films, or microcapsules. Sustained-release matrices include  
10 polylactides (U.S. Patent 3,773,919 and EP 58481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. et al., Biopolymers 22: 547-556 (1983)), poly (2-hydroxyethyl methacrylate) (R. Langer et al., J. Biomed. Mater. Res. 15: 167-277 (1981), and R. Langer, Chem. Tech. 12: 98-105 (1982)),  
15 ethylene vinyl acetate (R. Langer et al.) and poly-D- (-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also include liposomally entrapped polypeptides. Liposomes containing the polypeptide or other compound are prepared by well known methods (Epstein et al., Proc. Natl.  
20 Acad. Sci. USA 82: 3688-3692 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA 77: 4030-4034 (1980); EP 52322; EP 36676; EP 88046; EP 143949; EP 142641; Japanese Pat. Appl. 83-118008; U.S. Patent 4,485,045 and 4,544,545; and EP 102324). Ordinarily, the liposomes are of the small (about 200-800  
25 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal therapy.

For parenteral administration, in one embodiment, the polypeptide or other compound is formulated generally by  
30 mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation.

09886241-062101

For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to the polypeptide or other compound.

Generally, the formulations are prepared by contacting  
5 the polypeptide or other compound uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of  
10 the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier suitably contains minor amounts of additives  
15 such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic  
20 acid; low molecular weight (less than about ten residues) polypeptides, e. g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or  
25 arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates,  
30 poloxamers, or PEG.

The polypeptide or other compound is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the  
35 foregoing excipients, carriers, or stabilizers will result in

09886241.062101



the formation of polypeptide salts or salts of the other compounds.

Any polypeptide to be used for therapeutic administration should be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e. g., 0.2 micron membranes). Therapeutic polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Polypeptides ordinarily will be stored in unit or multi-dose containers, for example, sealed ampules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1 % (w/v) aqueous polypeptide solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized polypeptide using bacteriostatic Water-for-Injection.

#### 20 **Kits**

The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, reflecting approval by the agency of the manufacture, use or sale of the product for human administration.

#### 30 **Administration**

BSG polypeptides or polynucleotides or other compounds, preferably agonists or antagonists thereof of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

0986241-032101  
TOT290-14298360

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or  
5 intradermal routes among others.

The pharmaceutical compositions generally are administered in an amount effective for treatment or prophylaxis of a specific indication or indications. In general, the compositions are administered in an amount of at  
10 least about 10  $\mu\text{g/kg}$  body weight. However, it will be appreciated that optimum dosage will be determined by standard methods for each treatment modality and indication, taking into account the indication, its severity, route of administration, complicating conditions and the like.

15 It will be appreciated that conditions caused by a decrease in the standard or normal expression level of a BSG polypeptide in an individual can be treated by administering the BSG polypeptide of the present invention, preferably in the secreted form, or an agonist thereof. Thus, the invention  
20 also provides a method of treatment of an individual in need of an increased level of a BSG polypeptide comprising administering to such an individual a pharmaceutical composition comprising an amount of the BSG polypeptide or an agonist thereof to increase the activity level of the BSG  
25 polypeptide in such an individual. For example, a patient with decreased levels of a BSG polypeptide may receive a daily dose 0.1-100  $\mu\text{g/kg}$  of a BSG polypeptide or agonist thereof for six consecutive days. Preferably, if a BSG polypeptide is administered it is in the secreted form.

30 Compositions of the present invention can also be administered to treating increased levels of a BSG polypeptide. For example, antisense technology can be used to inhibit production of a BSG polypeptide of the present invention. This technology is one example of a method of  
35 decreasing levels of a polypeptide, preferably a secreted

0986241.062101  
TOT280.1428860

form, due to a variety of etiologies, such as cancer. A patient diagnosed with abnormally increased levels of a polypeptide can be administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 5 21 days. This treatment is preferably repeated after a 7-day rest period if the treatment was well tolerated. Compositions comprising an antagonist of a BSG polypeptide can also be administered to decrease levels of BSG in a patient.

#### **Gene therapy**

10 The BSG polynucleotides, polypeptides, agonists and antagonists that are polypeptides may be employed in accordance with the present invention by expression of such polypeptides *in vivo*, in treatment modalities often referred to as "gene therapy."

15 Thus, for example, cells from a patient may be engineered with a polynucleotide, such as a DNA or RNA, encoding a polypeptide *ex vivo*, and the engineered cells then can be provided to a patient to be treated with the polypeptide. For example, cells may be engineered *ex vivo* by 20 the use of a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention. Such methods are well-known in the art and their use in the present invention will be apparent from the teachings herein.

Similarly, cells may be engineered *in vivo* for 25 expression of a polypeptide *in vivo* by procedures known in the art. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed *supra*. The retroviral expression construct then may be isolated and introduced into 30 a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a patient for engineering cells

09806241.062101

in vivo and expression of the polypeptide in vivo. These and other methods for administering a polypeptide of the present invention would be apparent to those skilled in the art upon reading the instant application.

5           Retroviruses from which the retroviral plasmid vectors herein above mentioned may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human  
10 immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus. In one embodiment, the retroviral plasmid vector is derived from Moloney Murine Leukemia Virus.

Such vectors will include one or more promoters for  
15 expressing the polypeptide. The selection of a suitable promoter will be apparent to those skilled in the art from the teachings contained herein. However, examples of suitable promoters which may be employed include, but are not limited to, the retroviral LTR, the SV40 promoter, the human  
20 cytomegalovirus (CMV) promoter described in Miller et al., Biotechniques 7: 980-990 (1989), and eukaryotic cellular promoters such as the histone, RNA polymerase III, and beta-actin promoters. Other viral promoters which may be employed include, but are not limited to, adenovirus promoters,  
25 thymidine kinase (TK) promoters, and B19 parvovirus promoters. Additional promoters which may be used include respiratory syncytial virus (RSV) promoter, inducible promoters such as the MMT promoter, the metallothionein promoter, heat shock promoters, the albumin promoter, the ApoAI promoter, human  
30 globin promoters, viral thymidine kinase promoters such as the Herpes Simplex thymidine kinase promoter, retroviral LTRs, the beta-actin promoter, and human growth hormone promoters. The promoter also may be the native promoter which controls the gene encoding the polypeptide.

09886244-062101  
TOTAL 90-1129860

The nucleic acid sequence encoding the polypeptide of the present invention will be placed under the control of a suitable promoter.

In one embodiment, the retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, Y-2, Y-AM, PA12, T19-14X, VT-19-17-H2, YCRE, YCRIP, GP+E-86, GP+envAml2, and DAN cell lines as described in Miller, A., Human Gene Therapy 1: 5-14 (1990). The vector may be transduced into the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and  $\text{CaPO}_4$  precipitation. Alternatively, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host. The producer cell line will generate infectious retroviral vector particles which are inclusive of the nucleic acid sequence(s) encoding the polypeptides. Such retroviral vector particles then may be employed to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express the nucleic acid sequence(s) encoding the polypeptide. Eukaryotic cells which may be transduced include, but are not limited to, embryonic stem cells, embryonic carcinoma cells, as well as hematopoietic stem cells, hepatocytes, fibroblasts, myoblasts, keratinocytes, endothelial cells, and bronchial epithelial cells.

An exemplary method of gene therapy involves transplantation of fibroblasts which are capable of expressing a BSG polypeptide or an agonist or antagonist thereof onto a patient. Generally fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The

flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e. g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37°C for approximately one week. At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks. pMV-7 (Kirschmeier, P. T. et al., DNA, 7: 219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads. The cDNA encoding a BSG polypeptide of the present invention or an agonist or antagonist thereof can be amplified using PCR primers which correspond to their 5' and 3' end sequences respectively. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB 101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted. Amphotropic pA317 or GP+aml2 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now

09886241-063101  
TOTAL: 7428850

referred to as producer cells). Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced. The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

Alternatively, *in vivo* gene therapy methods can be used to treat BSG related disorders, diseases and conditions. Gene therapy methods relate to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to increase or decrease the expression of the polypeptide.

For example, a BSG polynucleotide of the present invention or a nucleic acid sequence encoding an agonist or antagonist thereto may be operatively linked to a promoter or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO 90/11092, WO 98/11779; U.S. Patents 5,693,622, 5,705,151, and 5,580,859; Tabata H. et al. (1997) Cardiovasc. Res. 35 (3): 470-479, Chao J et al. (1997) Pharmacol. Res. 35 (6): 517-522, Wolff J. A. (1997) Neuromuscul. Disord. 7 (5): 314-318, Schwartz B. et al. (1996) Gene Ther. 3 (5): 405-411,

Tsurumi Y. et al. (1996) Circulation 94 (12): 3281-3290 (incorporated herein by reference). The polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, 5 injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

The term "naked" polynucleotide, DNA or RNA, refers to 10 sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, polynucleotides may also be delivered in liposome 15 formulations (such as those taught in Felgner P. L. et al. (1995) Ann. NY Acad. Sci. 772: 126-139 and Abdallah B. et al. (1995) Biol. Cell 85 (1): 1-7) which can be prepared by methods well known to those skilled in the art.

The polynucleotide vector constructs used in the gene 20 therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies techniques, 25 one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for 30 periods of up to six months.

The polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, 35 kidney, gall bladder, stomach, intestine, testis, ovary,

09586241 "062101



uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred. The polynucleotide construct may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05  $\mu$ g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition,

naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template  
5 DNA for production of mRNA coding for polypeptide of the present invention is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are  
10 then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The template  
15 DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless  
20 steel clips.

After an appropriate incubation time (e. g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15  $\mu$ m cross-section of the individual quadriceps muscles is histochemically stained for protein  
25 expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT  
30 supernatants from injected and control mice.

The results of the above experimentation in mice can be use to extrapolate proper dosages and other treatment parameters in humans and other animals using naked DNA.

***Nonhuman Transgenic Animals***

The BSG polypeptides of the invention can also be expressed in nonhuman transgenic animals. Nonhuman animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, e. g., baboons, monkeys, and chimpanzees, may be used to generate transgenic animals. Any technique known in the art may be used to introduce the transgene (I. e., polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40: 691-698 (1994); Carver et al., Biotechnology (NY) 11: 1263-1270 (1993); Wright et al., Biotechnology (NY) 9: 830-834 (1991); and Hoppe et al., U.S. Patent 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82: 6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56: 313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol. Cell. Biol. 3: 1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, e. g., Ulmer et al., Science 259: 1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm mediated gene transfer (Lavitrano et al., Cell 57: 717-723 (1989)). For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115: 171-229 (1989), which is incorporated by reference herein in its entirety.

Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380: 64-66 (1996); Wilmut et al., Nature 385: 810813 (1997)).

05886241.062101

The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, i.e., mosaic or chimeric animals. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, e. g., head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89: 6232-6236 (1992)). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Science 265: 103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using

techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may  
5 also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the  
10 particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the  
15 transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis;  
20 crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of interest.

Transgenic animals of the invention have uses which  
25 include, but are not limited to, animal model systems useful in elaborating the biological function of BSG polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression of BSGs, and in screening for compounds effective in ameliorating such BSG associated  
30 conditions and/or disorders.

#### **Knock-Out Animals**

Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination (e. g., see Smithies  
35 et al., Nature 317: 230-234 (1985); Thomas & Capecchi, Cell

09886241-062101

51: 503512 (1987); Thompson et al., Cell 5: 313-321 (1989); each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional BSG polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous BSG polynucleotide sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (e. g., see Thomas & Capecchi 1987 and Thompson 1989, *supra*). This approach can also be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

In further embodiments of the invention, cells that are genetically engineered to express the BSG polypeptides of the invention, or alternatively, that are genetically engineered not to express the BSG polypeptides of the invention (e. g., knockouts) are administered to a patient *in vivo*. Such cells may be obtained from the patient or a MHC compatible donor and can include, but are not limited to, fibroblasts, bone marrow cells, blood cells (e. g., lymphocytes), adipocytes, muscle cells, and endothelial cells. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with

the polypeptides of the invention, e. g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of  
5 plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc.

The coding sequence of the BSG polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to  
10 achieve expression, and preferably secretion, of the BSG polypeptides of the invention. The engineered cells which express and preferably secrete the BSG polypeptides of the invention can be introduced into the patient systemically, e.g., in the circulation, or intraperitoneally.

15 Alternatively, the cells can be incorporated into a matrix and implanted in the body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft or genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft (see, for example, U.S.  
20 Patent 5,399,349 and U.S. Patent 5,460,959 each of which is incorporated by reference herein in its entirety).

When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the development of a host  
25 immune response against the introduced cells. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

30 Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of BSG polypeptides of the present invention, studying conditions and/or disorders associated with aberrant BSG expression, and

09886241.062101

in screening for compounds effective in ameliorating such BSG associated conditions and/or disorders.

#### EXAMPLE

5           The present invention is further described by the following example. The example is provided solely to illustrate the invention by reference to specific embodiments. This exemplification, while illustrating certain aspects of the invention, does not portray the limitations or  
10 circumscribe the scope of the disclosed invention.

          All examples outlined here were carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. Routine molecular biology techniques of the following  
15 example can be carried out as described in standard laboratory manuals, such as Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

#### 20   Semi-quantitative Polymerase Chain Reaction (SQ-PCR):

          Semi-quantitative Polymerase Chain Reaction (SQ-PCR) is a method that utilizes end point PCR on serial dilutions of cDNA samples in order to determine relative expression patterns of genes of interest in multiple samples. Using  
25 random hexamer primed Reverse Transcription (RT) cDNA panels are created from total RNA samples. Gene specific primers are then used to amplify fragments using Polymerase Chain Reaction (PCR) technology from four 10x serial cDNA dilutions in duplicate. Relative expression levels of 0, 1, 10, 100 and  
30 1000 are used to evaluate gene expression. A positive reaction in the most dilute sample indicates the highest relative expression value. This is determined by analysis of the sample reactions on a 2-4% agarose gel. The tissue samples used include 12 normal, 12 cancer and 6 pairs of  
35 tissue specific cancer and matching samples. Leads evaluated

09886241.062101





Table 2 shows absolute numbers which are relative levels of expression of sqmam040 (SEQ ID NO:1) in 12 cancer samples from 12 different tissues. Using Polymerase Chain Reaction (PCR) technology expression levels were analyzed from four 10x 5 serial cDNA dilutions in duplicate. Relative expression levels of 0, 1, 10, 100 and 1000 are used to evaluate gene expression. A positive reaction in the most dilute sample indicates the highest relative expression value

Table 2:

10	<b>TISSUE</b>	<b>CANCER</b>
	bladder	100
	breast	100
	colon	10
	kidney	10
15	liver	10
	lung	100
	ovary	100
	pancreas	10
	prostate	10
20	stomach	10
	testes	100
	uterus	1000

Relative levels of expression in Table 2 show that sqmam040 25 (SEQ ID NO:1) is expressed in all the carcinomas at different degrees.

Table 3 shows absolute numbers which are relative levels of expression of sqmam040 (SEQ ID NO:1) in 6 mammary gland cancer matching samples. A matching pair is formed by mRNA 30 from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual. Using Polymerase Chain Reaction (PCR) technology expression levels were analyzed from four 10x serial cDNA dilutions in duplicate. Relative expression levels of 0, 1, 35 10, 100 and 1000 are used to evaluate gene expression. A positive reaction in the most dilute sample indicates the highest relative expression value.

Table 3:

	SAMPLE ID	TISSUE	CANCER	NORMAL ADJACENT TISSUE
	S99522A/B	mammary gland 1	1	10
5	4005724A2/B3	mammary gland 2	100	10
	4005599A4/B2	mammary gland 3	1	1
	4005629A2/B2	mammary gland 4	1	1000
	S9822245A/B	mammary gland 5	1000	10
10	S9819997A/B	mammary gland 6	100	1000

Relative levels of expression in Table 3 shows that sqmam040 (SEQ ID NO:1) is expressed in all six mammary gland matching samples. This assay shows that sqmam040 (SEQ ID NO:1) is upregulated in 2 (33%) matching samples analyzed, downregulated in 3 (50%), and shows no difference in one matching sample.

CHROMOSOME LOCALIZATION: Blast result against the human genome database, showed no significant homology to chromosome sequences.

20

**Sqmam041; SEQ ID NO:2**

Semi-quantitative PCR was done using the following primers:

sqmam041 forward:

25 5' CCTGTGACACCTGTAGCCTAATAAA 3' (SEQ ID NO:9)

sqmam041 reverse:

5' TGACGCTCTGCTGTAAGAAAAA 3' (SEQ ID NO:10)

Table 4 shows absolute numbers which are relative levels of expression of sqmam041 (SEQ ID NO:2) in 12 normal samples from 12 different tissues. These RNA samples are from single individual or are commercially available pools, originated by pooling samples of a particular tissue from different individuals. Using Polymerase Chain Reaction (PCR) technology expression levels were analyzed from four 10x serial cDNA dilutions in duplicate. Relative expression levels of 0, 1, 10, 100 and 1000 are used to evaluate gene expression. A

positive reaction in the most dilute sample indicates the highest relative expression value.

Table 4:

	TISSUE	NORMAL
5	Breast	100
	Colon	100
	Endometrium	1000
	Kidney	100
	Liver	10
10	Lung	100
	Ovary	1
	Prostate	100
	Small Intestine	1000
	Stomach	100
15	Testis	10
	Uterus	100

Relative levels of expression in Table 4 show that sqmam041 (SEQ ID NO:2) is expressed in all the normal samples tested.

Table 5 shows absolute numbers which are relative levels of expression of sqmam041 (SEQ ID NO:2) in 12 cancer samples from 12 different tissues. Using Polymerase Chain Reaction (PCR) technology expression levels were analyzed from four 10x serial cDNA dilutions in duplicate. Relative expression levels of 0, 1, 10, 100 and 1000 are used to evaluate gene expression. A positive reaction in the most dilute sample indicates the highest relative expression value.

Table 5:

	TISSUE	CANCER
30	bladder	10
	breast	100
	colon	10
	kidney	0
	liver	0
35	lung	1
	ovary	1000
	pancreas	10
	prostate	0
	stomach	10
40	testes	1000
	uterus	100

Relative levels of expression in Table 5 show that sqmam041 (SEQ ID NO:5) is not expressed in kidney, liver and prostate carcinomas, and expressed at different levels in the rest of the tissues.

5 Table 6 shows absolute numbers which are relative levels of expression of sqmam041 (SEQ ID NO:2) in 6 mammary gland cancer matching samples. A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same  
10 individual. Using Polymerase Chain Reaction (PCR) technology expression levels were analyzed from four 10x serial cDNA dilutions in duplicate. Relative expression levels of 0, 1, 10, 100 and 1000 are used to evaluate gene expression. A  
15 positive reaction in the most dilute sample indicates the highest relative expression value.

Table 6:

	SAMPLE ID	TISSUE	CANCER	NORMAL ADJACENT
				TISSUE
	S99522A/B	mammary gland 1	100	10
20	4005724A2/B3	mammary gland 2	100	10
	4005599A4/B2	mammary gland 3	10	10
	4005629A2/B2	mammary gland 4	100	100
	S9822245A/B	mammary gland 5	10	100
25	S9819997A/B	mammary gland 6	10	10

Relative levels of expression in Table 6 shows that sqmam041 (SEQ ID NO:2) is expressed in all the matching mammary gland samples tested. This assay shows that sqmam041 (SEQ ID NO:2) is upregulated in 2 (33%) of the matching samples analyzed;  
30 downregulated in 2 (33%) and shows no differential expression in another 2 matching samples.

CHROMOSOME LOCALIZATION: Blast result against the human genome database, showed no significant homology to chromosome  
35 sequences.

**Sqmam042 (SEQ ID NO:4)**

Semi-quantitative PCR was done using the following primers:

sqmam042 forward:

5' GCCTTACAGAGACTGGAAAAGAA 3' (SEQ ID NO:11)

5 sqmam042 reverse:

5' TCAGCGAAGGTGGTGTGTAG 3' (SEQ ID NO:12)

Table 7 shows absolute numbers which are relative levels of expression of sqmam042 (SEQ ID NO:4) in 12 normal samples from 12 different tissues. These RNA samples are from single individual or are commercially available pools, originated by pooling samples of a particular tissue from different individuals. Using Polymerase Chain Reaction (PCR) technology expression levels were analyzed from four 10x serial cDNA dilutions in duplicate. Relative expression levels of 0, 1, 10, 100 and 1000 are used to evaluate gene expression. A positive reaction in the most dilute sample indicates the highest relative expression value.

**Table 7:**

	<b>TISSUE</b>	<b>NORMAL</b>
20	Breast	10
	Colon	1000
	Endometrium	10
	Kidney	10
	✓ Liver	0
25	Lung	10
	Ovary	10
	Prostate	10
	Small Intestine	100
	Stomach	10
30	Testis	100
	Uterus	100

Relative levels of expression in Table 7 show that except for normal liver, sqmam042 (SEQ ID NO:4) is expressed in all the other normal tissues.

Table 8 shows absolute numbers which are relative levels of expression of sqmam042 (SEQ ID NO:4) in 12 cancer samples from 12 different tissues. Using Polymerase Chain Reaction (PCR)

technology expression levels were analyzed from four 10x serial cDNA dilutions in duplicate. Relative expression levels of 0, 1, 10, 100 and 1000 are used to evaluate gene expression. A positive reaction in the most dilute sample 5 indicates the highest relative expression value.

Table 8:

	TISSUE	CANCER
	bladder	10
	breast	1
10	colon	10
	kidney	10
	liver	1
	lung	10
	ovary	1000
15	pancreas	10
	prostate	10
	stomach	10
	testes	10
20	uterus	10

Relative levels of expression in Table 8 show that sqmam042 (SEQ ID NO:4) is expressed in all the carcinomas tested.

Table 9 shows absolute numbers which are relative levels of expression of sqmam042 (SEQ ID NO:4) in 6 mammary gland cancer matching samples. A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual. Using Polymerase Chain Reaction (PCR) technology expression levels were analyzed from four 10x serial cDNA dilutions in duplicate. Relative expression levels of 0, 1, 10, 100 and 1000 are used to evaluate gene expression. A positive reaction in the most dilute sample indicates the highest relative expression value.

Table 9:

	SAMPLE ID	TISSUE	CANCER	NORMAL ADJACENT TISSUE
35	S99522A/B	mammary gland 1	1000	1
	4005724A2/B3	mammary gland 2	1000	1000
	4005599A4/B2	mammary gland 3	1000	10
40	4005629A2/B2	mammary gland 4	10	10
	S9822245A/B	mammary gland 5	1000	1000

S9819997A/B      mammary gland 6      1000      1000

Relative levels of expression in Table 9 shows that sqmam042 (SEQ ID NO:4) is expressed in all six matching mammary gland 5 samples. This assay shows that sqmam042 is upregulated in 2 (33%) of the matching samples analyzed, ~~and the other 4 (66%) matching samples did not show differential expression between~~ normal and cancer.

- 10 CHROMOSOME LOCALIZATION: Blast result against the human genome database, showed that the sequence listed below for sqmam042 has 95-100% identity with a sequence in chromosome 17.

0988241.062401  
"F4238850"